

Today's Date: _____ Needed by: _____ Name: _____ Phone: _____

Patient Demographic: (Provide the following or attached demographic sheet) Patient Name: _____ Address: _____ City, State, Zip: _____ Home Phone: _____ Alt. Phone: _____ SS#: _____ Date Birth: _____ Gender: _____ Height: _____ Weight: _____ BSA: _____ m2	Prescriber: (Provide as much information as possible) Prescriber's Name: _____ Group/Hospital: _____ Specialty: _____ NPI #: _____ DEA: _____ Address: _____ City, State, Zip: _____ Phone: _____ Fax: _____ Alt. Contact Name: _____ Phone: _____ Position: _____
---	--

Insurance Information (Please copy and attach the front and back of the Insurance Card):
 Primary Insurance Name: _____ ID# _____ BIN#: _____ Group#: _____ PCN#: _____ Phone: _____
 No Insurance Patient will pay out of pocket Enroll in Manufacturer's Patient Assistance Program

Medication Delivery to (Choose Only One):
 Patient Address First Fill Physician's Office, Refill to Patient Address Patient will pick up at Pharmacy

Diagnosis (ICD- 10 code):
 B17.10 Acute Hepatitis C without hepatic come B17.11 Acute Hepatitis C with hepatic come B18.2 Chronic Hepatitis C
 B19.20 Unspecified Viral Hepatitis C without hepatic coma B20 HIV T86.40 Liver Transplant Other: _____

Injection Training & Educational Needs: Specialty Pharmacy Injection Training Requested Manufacturer's Patient Assistance Program Enrollment Requested **OR** (Please Choose only one) Physician's Office already trained Patient Patient is already independently Injecting

Patient Evaluation:
 Height: _____ in/cm Weight: _____ kg/lbs. Allergies: _____
 HCV Genotype 1a 1b 1 2 3 4 5 6 AND No Cirrhosis Compensated Cirrhosis Uncompensated Cirrhosis
 Is patient: Naïve Partial Responder Non-Responder Relapser; Last Date of Therapy: _____ Product Names: _____
 Is patient currently on Hepatitis C Virus (HCV) therapy? Yes No; If Yes, Therapy Start Date: _____ Product Names: _____

MEDICATION	STRENGTH	DIRECTION	Qty. SUPPLY	REFILLS
<input type="checkbox"/> Daklinza (daclatasvir)	<input type="checkbox"/> 60mg tablets <input type="checkbox"/> 30mg tablets <input type="checkbox"/> 90mg tablets	<input type="checkbox"/> Take one 60mg tablet orally once a day. <input type="checkbox"/> Take one 90mg tablet orally once a day. <input type="checkbox"/> Other: _____	28 day	<input type="checkbox"/> 12 weeks
<input type="checkbox"/> Epclusa (sofosbuvir/velpatasvir)	<input type="checkbox"/> One tablet contains 400 mg sofosbuvir and 100 mg velpatasvir.	One tablet orally once a day with or without food (Patients with decompensated cirrhosis, add Ribavirin)	28 day	<input type="checkbox"/> 12 weeks <input type="checkbox"/> ___ weeks
<input type="checkbox"/> Harvoni (ledipasvir/sofosbuvir)	<input type="checkbox"/> Fixed-dose combination tablet of 90mg of ledipasvir/400mg of sofosbuvir.	Take orally once daily with or without food (Do not take within 4 hours of antacids)	28 day	<input type="checkbox"/> 8 weeks <input type="checkbox"/> 12 weeks <input type="checkbox"/> 24 weeks
<input type="checkbox"/> Olysio (simeprevir)	<input type="checkbox"/> 150 mg caps	Take one 150mg cap orally once a day.	28 day	<input type="checkbox"/> 12 weeks <input type="checkbox"/> 24 weeks
<input type="checkbox"/> Ribavirin	<input type="checkbox"/> 200 mg tabs <input type="checkbox"/> 200mg caps	Take ___ tabs/caps PO QAM and ___ tabs/caps PO QPM to equal a total of ___mg/day		
<input type="checkbox"/> Risbasphere RIBA-PAK	<input type="checkbox"/> 600/600 mg tabs <input type="checkbox"/> 600/400mg <input type="checkbox"/> 400/400 mg tabs <input type="checkbox"/> 200/400mg	Take ___ mg PO QAM and ___ QPM to equal a total of ___mg/day		
<input type="checkbox"/> Sovaldi (sofosbuvir)	<input type="checkbox"/> 400 mg tabs	Take one 400mg tab orally once a day	28 day	
<input type="checkbox"/> Technivie (ombitasvir/paritaprevir/ritonavir)	Fixed dose combination tablet of ombitasvir/paritaprevir/ritonavir (12.5/75/50 mg)	Take two tablets (ombitasvir, paritaprevir, ritonavir) once daily in the morning.	28 day	<input type="checkbox"/> 12 weeks
<input type="checkbox"/> Victrelis (boceprevir)	<input type="checkbox"/> 200 mg caps	Take 800mg orally 3 times a day every 7-9 hours with food. (Begin after week 4 of PEGylated interferon therapy)	28 day	
<input type="checkbox"/> Viekira Pak (ombitasvir/paritaprevir/ritonavir tablets and dasabuvir tablets)	Viekira Pak: ombitasvir/paritaprevir/ritonavir 12.5/75/50 mg and dasabuvir 250 mg copackaged	Take two pink tablets (ombitasvir, paritaprevir, ritonavir) once daily (morning) and one beige tablet (dasabuvir) twice daily (morning and evening) with meals.	28 day	<input type="checkbox"/> 12 weeks <input type="checkbox"/> 24 weeks
<input type="checkbox"/> Viekira XR (dasabuvir/ombitasvir/paritaprevir/ritonavir)	Extended release tablets 200mg dasabuvir, 8.33 mg ombitasvir, 50mg paritaprevir and 33.33 mg ritonavir	Take 3 tablets once daily with a meal <input type="checkbox"/> Add Ribavirin	28 day	<input type="checkbox"/> 12 weeks <input type="checkbox"/> 24 weeks
<input type="checkbox"/> Zepatir (elbasvir/grazoprevir)	<input type="checkbox"/> One tablet contains 50 mg elbasvir and 100 mg grazoprevir	One tablet orally once a day with or without food <input type="checkbox"/> Add Ribavirin	28 day	<input type="checkbox"/> 12 weeks <input type="checkbox"/> 16 weeks
<input type="checkbox"/> Other				

By signing this form and utilizing our services, you are authorizing SMP and its employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

_____ / _____ / _____
 Prescriber Signature Date